



**DuPont Crop
Protection**

SUPPLEMENTAL LABELING

**DUPONT™ CORAGEN®
INSECT CONTROL WITH ACTIVE
INGREDIENT RYNAXYPYR®
FOR USE ON VARIOUS
VEGETABLE CROPS**

DUPONT™ CORAGEN® INSECT CONTROL

EPA Reg. No. 352-729

FOR FOLIAR INSECT CONTROL ON ARTICHOKE, ASPARAGUS, CORN (FIELD, POP, SEED, SWEET), LEGUME VEGETABLES, OKRA, STRAWBERRY, SUGARCANE, TOBACCO & TUBEROUS AND CORM VEGETABLES

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

CORAGEN® is a suspension concentrate that can be applied as a foliar spray to control listed insects. CORAGEN® is mixed with water for application.

RESISTANCE MANAGEMENT

For resistance management, CORAGEN® is a Group 28 Insecticide. Repeated and exclusive use of CORAGEN® (chlorantraniliprole) or other Group 28 insecticide belonging to the anthranilic diamide class of chemistry may lead to the buildup of resistant strains of insects in some crops.

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, this product may be used as part of a resistance management strategy established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode-of-action classes of insecticides on succeeding generations and the most susceptible life stage. Consult your local or state agricultural authorities for details.

Unless directed otherwise in the specific crop/pest sections of this label, the best practices are to follow these instructions to delay the development of insecticide resistance: Avoid using the same mode of action (same insecticide group) on consecutive generations of insect pests. Make no more than 2 applications of DuPont™ CORAGEN® (chlorantraniliprole) or other Group 28 products per generation to the same insect species on a crop. Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide). Make no more than 2 successive applications within a 30

day period to the same insect species on a crop. The following application to the target pest(s) must be with an effective product with a different mode of action. See Federal Section 3 label accompanying product for full resistance management instructions.

APPLICATION

Apply at the specified rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Apply follow-up treatments of CORAGEN®, as specified in the tables on page 2 of this label, to keep pest populations within threshold limits. Refer to the Resistance Management section of this label for further guidance on follow-up treatments. See individual crop sections of this label for specific minimum spray intervals.

Use sufficient water to obtain thorough, uniform coverage.

IMPORTANT

Before using CORAGEN(R), read and follow all applicable directions: restrictions; and precautions on the EPA-registered label. This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

© 2010 E. I. du Pont de Nemours and Company, Crop Protection, 1007 N. Market St., Wilmington, Delaware 19898

Crop	Application Method	Target Pest	DUPONT™ CORAGEN® RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb. ai per acre	fluid ounces product per acre		
Artichoke	FOLIAR	Artichoke plume moth	0.045-0.098	3.5 - 7.5	3	4
	<p>Make no more than 4 applications per acre per crop. Do not apply more often than each 14 days. Apply no more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop. Make applications between bud formation and harvest of an individual fruit. Apply in a minimum of 10 gallons water per acre by air and 50 - 200 gallons of water per acre by ground (use sufficient water to obtain thorough coverage without excessive runoff).</p>					
Asparagus	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045-0.065	3.5 - 5.0	1	4
	<p>Make no more than 4 applications per acre per crop. Minimum interval between treatments is 3 days. Do not use an adjuvant with applications of CORAGEN®. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.</p>					
Corn (field) Corn (pop)	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm	0.045-0.065	3.5 - 5.0	14	4
	<p>Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.</p>					
Corn (sweet) Corn (seed)	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm	0.045-0.065	3.5 - 5.0	1	4
	<p>Make no more than 4 applications per acre per crop. Minimum interval between treatments is 1 day. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.</p>					

Crop	Application Method	Target Pest	DUPONT™ CORAGEN® RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb. ai per acre	fluid ounces product per acre		
Legume vegetables except soybean (Succulent or Dried): Bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (Vigna) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (Pisum) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; sword bean	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm	0.045-0.065	3.5 - 5.0	1	4
<p>Make no more than 4 applications per acre per crop. Minimum interval between treatments is 3 days. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.</p>						
Okra	FOLIAR	Beet armyworm Corn earworm	0.045-0.065	3.5 - 5.0	1	4
<p>Make no more than 4 applications per acre per crop. Minimum interval between treatments is 5 days. Do not use an adjuvant with applications of CORAGEN®. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.</p>						

Crop	Application Method	Target Pest	DUPONT™ CORAGEN® RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb. ai per acre	fluid ounces product per acre		
Strawberry	FOLIAR	Beet armyworm Cabbage looper Corn earworm Japanese beetle (adult)	0.045 - 0.065	3.5 - 5.0	1	4
	Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.					
Sugarcane	FOLIAR	Sugarcane borer	0.045-0.065	3.5 - 5.0	14	4
	Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.					
Tobacco	FOLIAR	Split worm (potato tuberworm) Tobacco budworm Tomato hornworm Tobacco hornworm	0.045-0.098	3.5 - 7.5	1	4
	Make no more than 4 applications per acre per crop. Minimum interval between treatments is 3 days. Do not use an adjuvant with applications of CORAGEN®. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.					
Tuberous and Corm Vegetables (except potato): Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; sweet potato; tanier; turmeric; yam bean; yam, true	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045-0.065	3.5 - 5.0	14	4
	Make no more than 4 applications per acre per crop. Minimum interval between treatments is 5 days. Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.					